arXiv:1409.2630v1 [math.FA] 9 Sep 2014

J-class sequences of linear operators

M.R. Azimi

Department of Mathematics, Faculty of Science, University of Maragheh, 55181-83111 Maragheh, Iran.

Abstract

In this paper we first introduce the extended limit set $J_{\{T^n\}}(x)$ for a sequence of bounded linear operators $\{T_n\}_{n=1}^{\infty}$ on a separable Banach space X . Then we study the dynamics of sequence of linear operators by using the extended limit set. It is shown that the extended limit set is strongly related to the topologically transitive of a sequence of linear operators. Finally we show that a sequence of operators $\{T_n\}_{n=1}^{\infty} \subseteq \mathcal{B}(X)$ is hypercyclic if and only if there exists a cyclic vector $x \in X$ such that $J_{\{T^n\}}(x) = X$.

Keywords: sequences of operators, hypercyclic operators, J-class operators,
2010 MSC: Primary 47A16 Secondary 37B99

1. Introduction and Preliminaries

Let X be a separable Banach space over the field of complex numbers $\mathbb{C}$ and let $\mathcal{B}(X)$ be the algebra of all bounded linear operators on X. An operator $T \in \mathcal{B}(X)$ is called *hypercyclic* if there exists a vector $x \in X$ whose orbit under T, $Orb(T, x) = \{T^n x : n \in \mathbb{N}\}$, is dense in X. Such a vector is called *hypercyclic vector* for T. An operator $T \in \mathcal{B}(X)$ is called *topologically transitive* if for any pair U, V of nonempty open subsets of X, there exits $n \in \mathbb{N}$ such that

$$T^n(U) \cap V \neq \emptyset.$$

Over last two decades many useful criteria and results have been presented for the hypercyclicity of linear operators which may find in [10], [3], [15], [12] and

Email address: mhr.azimi@maragheh.ac.ir (M.R. Azimi)

the references therein. It is well known and easy to prove that an operator $T \in \mathcal{B}(X)$ is hypercyclic if and only if T is topologically transitive (see [3]).
In recent years the dynamics of linear operators has been widely studied in terms of their orbits by many authors. The survey articles [8], [5], [5], [12], [14] and the book [3] are good references in this subject. In [8] G. Costakis and A. Manoussos have studied the dynamics of linear operators by making use of extended limit sets instead of their orbits. Especially a result due to P. Bourdon and N. Feldman ([6]) has been generalized by them as follows.

Theorem 1.1. *([8]) If x is a cyclic vector for an operator $T \in \mathcal{B}(X)$ and the set $J_T(x)$ has non-empty interior then $J_T(x) = X$ and so T is hypercyclic.*

They have also posed several open problems which one of them was answered in [2]. In particular they have studied J-class weighted shifts in [7]. It should be mentioned that H. Salas is the first mathematician who characterized the hypercyclicity of weighted shifts in [16]. Recently Q. Menet [13] generalized the results of F. León-Saavedra and V. Müller [11] and obtained a new criterion for characterizing of weighted shifts with hypercyclic subspaces.

Let us define the notion of extended limit set. Let $T \in \mathcal{B}(X)$ and $x \in X$. The *extended limit set* of x, denoted by $J_T(x)$ consists of those vectors y in X if there exist a strictly increasing sequence of positive integers $\{k_n\}$ and a sequence $\{x_n\} \subseteq X$ such that $x_n \to x$ and $T^{k_n} x_n \to y$. An operator T is called J-class operator whenever $J_T(x) = X$ for some non-zero vector $x \in X$.

Actually the extended limit set localizes the notion of hypercyclicity. For more details the reader is referred to [8].

The goal of this paper is to investigate the dynamics of sequences of linear operators using the localized hypercyclicity notion. For this we start by the definition of hypercyclic and topologically transitive sequences of linear operators.

Definition 1.1. A sequence of operators $\{T_n\}_{n=1}^{\infty} \subseteq \mathcal{B}(X)$ is called *hypercyclic* if there exists a vector $x \in X$ such that the set $\{T_n x : n \in \mathbb{N}\}$ is dense in X. Such a vector is called hypercyclic vector for the sequence of operators $\{T_n\}_{n=1}^{\infty}$.

We say that an operator $T : X \to X$ is hypercyclic if the sequence of its iterates $\{T^n\}_{n=1}^{\infty}$ is hypercyclic.

Definition 1.2. A sequence of operators $\{T_n\}_{n=1}^{\infty} \subseteq \mathcal{B}(X)$ is called *topologically transitive* if for any pair U, V of nonempty open subsets of X, there exits $n \in \mathbb{N}$ such that

$$T_n(U) \cap V \neq \emptyset.$$

The sequence of linear operators $\{T_n\}_{n=1}^{\infty}$ is hypercyclic if and only if it is topologically transitive([9]). The hypercyclicity criterion for sequences of operators has been studied in [4]. In [11], F. León-Saavedra and V. Müller have studied the various conditions and criteria that imply the hypercyclicity of $\{T_n\}_{n=1}^{\infty}$.

In this paper we first introduce the extended limit set for a sequence of bounded linear operators on a separable Banach space X analogous to the single operator case. Then we study the dynamics of sequence of linear operators by using the extended limit set. It is turn out that the extended limit set is strongly related to the hypercyclicity of a sequence of linear operators. Indeed we show that a sequence of operators $\{T_n\}_{n=1}^{\infty} \subseteq \mathcal{B}(X)$ is hypercyclic if and only if there exists a cyclic vector $x \in X$ such that $J_{\{T^n\}}(x) = X$. Recall that a vector x is *cyclic* for $\{T_n\}_{n=1}^{\infty}$ if the linear span of the set $\{T_n x : n \in \mathbb{N}\}$ is dense in X.

Definition 1.3. Let $\{T_n\}_{n=1}^{\infty} \subseteq \mathcal{B}(X)$ and $x \in X$. The *extended limit set* of x under $\{T_n\}_{n=1}^{\infty}$, denoted by $J_{\{T_n\}}(x)$ is the collection of all vectors y in X if there exist a strictly increasing sequence of positive integers $\{k_n\}$ and a sequence $\{x_n\} \subseteq X$ such that $x_n \to x$ and $T_{k_n} x_n \to y$. A sequence of bounded linear operators $\{T_n\}_{n=1}^{\infty}$ is called J-class sequence of linear operators whenever $J_{\{T_n\}}(x) = X$ for some non-zero vector $x \in X$.

By comparing two following definitions

$$J_T(x) = \{y \in X : \text{there exist a strictly increasing sequence of positive integers } \{k_n\} \text{ and a sequence } \{x_n\} \subseteq X \text{ such that } x_n \to x \text{ and } T^{k_n} x_n \to y\}.$$

and

$$J_{\{T_n\}}(x) = \{y \in X : \text{there exist a strictly increasing sequence of positive integers } \{k_n\} \text{ and a sequence } \{x_n\} \subseteq X \text{ such that } x_n \to x \text{ and } T_{k_n}x_n \to y\}.$$

it is found out that $J_T(x) = J_{\{T^n\}}(x)$ for every $x \in X$.

2. Main Results

In this section we review some basic properties of the extended limit sets. Afterwards we characterize the hypercyclicity of $\{T_n\} \subseteq \mathcal{B}(X)$ based upon these sets.

Lemma 2.1. *If $\{T_n\}$ is hypercyclic then $J_{\{T_n\}}(x) = X$ for every $x \in X$.*

Proof. Let $x, y \in X$ be arbitrary. Remind that the set of all hypercyclic vectors for $\{T_n\}$ is dense in X, see [9]. Hence there exists a sequence of hypercyclic vectors $\{x_n\}$ such that $x_n \to x$. But in view of the hypercyclicity definition, the set $\{T_n x : n \in \mathbb{N}\}$ being dense in X, one may find a strictly increasing sequence of positive integers $\{k_n\}$ such that $T_{k_n}x_n \to y$. This means that $y \in J_{\{T_n\}}(x)$ and so $J_{\{T_n\}}(x) = X$. □

Remark 2.1. *Note that an equivalent definition of $J_{\{T_n\}}(x)$ can be stated in the following.*

$$J_{\{T_n\}}(x) = \{y \in X : \textit{for every pair neighborhoods } U, V \textit{ of } x, y \textit{ respectively, there exists a positive integer } n \textit{ such that } T_nU \cap V \neq \emptyset\}.$$

Observe now that $\{T_n\}$ is topologically transitive if and only if $J_{\{T_n\}}(x) = X$ for every $x \in X$.

Lemma 2.2. *Let $x_n \to x$ and $y_n \to y$ for some $x, y \in X$. If $y_k \in J_{\{T_n\}}(x_k)$ for every $k = 1, 2, 3, ...$, then $y \in J_{\{T_n\}}(x)$.*

Proof. The proof is followed same as the single operator case stated in [7]. By our hypothesis there exists a positive integer k_1 such that $\|x_{k_1} - x\| < \frac{1}{2}$

and $\|y_{k_1} - y\| < \frac{1}{2}$. Since $y_{k_1} y \in J_{\{T_n\}}(x_{k_1})$ there exist a positive integer l_1 and $z_1 \in X$ such that $\|z_1 - x_{k_1}\| < \frac{1}{2}$ and $\|T_{l_1} z_1 - y_{k_1} y\| < \frac{1}{2}$. Hence $\|z_1 - x\| < 1$ and $\|T_{l_1} z_1 - y\| < 1$ which completes the firs step of the induction. By proceeding inductively we now construct a strictly increasing sequence of positive integers $\{l_n\}$ and a sequence of vectors $\{l_n\}$ in X such that $\|z_n - x\| < \frac{1}{n}$ and $\|T_{l_n} z_n - y\| < \frac{1}{n}$. Therefore $y \in J_{\{T_n\}}(x)$. □

Lemma 2.3. *Let $\{T_n\} \subseteq \mathcal{B}(X)$ such that $T_n \to T$ uniformly on X for some $T \in \mathcal{B}(X)$. If $T_n y \in J_{\{T_n\}}(T_n x)$ for every $n = 1, 2, 3, ...$ and for arbitrary $x, y \in X$, then $Ty \in J_{\{T_n\}}(Tx)$.*

Proof. For an arbitrary $\epsilon > 0$ there exists a positive integer k_1 such that $\|T_{k_1} - T\| < \frac{\epsilon}{2}$. Since $T_{k_1} y \in J_{\{T_n\}}(T_{k_1} x)$ there exist a positive integer l_1 and $z_1 \in X$ such that $\|z_1 - T_{k_1} x\| < \frac{\epsilon}{2}$ and $\|T_{l_1} z_1 - T_{k_1} y\| < \frac{\epsilon}{2}$. Hence $\|z_1 - Tx\| < \epsilon$ and $\|T_{l_1} z_1 - Ty\| < \epsilon$ i.e., $Ty \in J_{\{T_n\}}(Tx)$. □

Definition 2.1. Let $\{T_n\}_{n=1}^{\infty} \subseteq \mathcal{B}(X)$ and $x \in X$. The *limit set* of x under $\{T_n\}_{n=1}^{\infty}$, denoted by $L_{\{T_n\}}(x)$ is defined as follows

$$L_{\{T_n\}}(x) = \{y \in X : \text{there exist a strictly increasing sequence of positive integers } \{k_n\},\ T_{k_n} x \to y\}.$$

In case $\{T_n\}_{n=1}^{\infty}$ is sequence of invertible linear operators the sets $L^{+}_{\{T_n\}}(x)$, $J^{+}_{\{T_n\}}(x)$ ($L^{-}_{\{T_n\}}(x)$, $J^{-}_{\{T_n\}}(x)$) denote the limit set and extended limit set of x under $\{T_n\}_{n=1}^{\infty}$ ($\{T_n^{-1}\}_{n=1}^{\infty}$).

Lemma 2.4. *Let $\{T_n\}$ be a sequence of mutually commuting operators in $\mathcal{B}(X)$. For all $x \in X$, the sets $J_{\{T_n\}}(x)$ and $L_{\{T_n\}}(x)$ are closed. Moreover for each $T_m \in \mathcal{B}(X)$ we have $T_m J_{\{T_n\}}(x) \subseteq J_{\{T_n\}}(T_m x)$ and $T_m L_{\{T_n\}}(x) \subseteq L_{\{T_n\}}(T_m x)$.*

Proof. Take $y \in J_{\{T_n\}}(x)$. There exist a strictly increasing sequence of positive integers $\{k_n\}$ and a sequence $\{x_n\}$ in X such that $x_n \to x$ and $T_{k_n} x_n \to y$. Then $T_{k_n} T_m x_n = T_m T_{k_n}(x) \to T_m y$ and so $T_m y \in J_{\{T_n\}}(T_m x)$ since $T_m x_n \to T_m x$. □

Lemma 2.5. *Let $\{T_n\}$ be a commuting sequence of invertible operators. Then $T_n^{-1}J_{\{T_n\}}(x) = J_{\{T_n\}}(T_nx)$ for every $x \in X$.*

Proof. By Lemma 2.4, we have $J_{\{T_n\}}(x) \subseteq T_n^{-1}J_{\{T_n\}}(T_nx)$. Now pick $y \in T_n^{-1}J_{\{T_n\}}(x)$. There are a strictly increasing sequence of positive integers $\{k_i\}$ and a sequence $\{x_i\}$ in X such that $x_i \to T_nx$ and $T_{k_i}x_i \to T_ny$ as $i \to \infty$. Hence $T_n^{-1}x_i \to x$ and $T_{k_i}T_n^{-1}x_i = T_n^{-1}T_{k_i}x_i \to y$. That is $y \in J_{\{T_n\}}(x)$. □

Proposition 2.1. *Let $\{T_n\}$ be a sequence of invertible operators. Then $y \in J^+_{\{T_n\}}(x)$ if and only if $x \in J^-_{\{T_n\}}(y)$ for every $x, y \in X$.*

Proof. Choose $y \in J^+_{\{T_n\}}(x)$. There are a strictly increasing sequence of positive integers $\{k_n\}$ and a sequence $\{x_n\}$ in X such that $x_n \to x$ and $T_{k_n}x_n \to y$. Then $T_{k_n}^{-1}T_{k_n}x_n = x_n \to x$ and so $x \in J^-_{\{T_n\}}(y)$. The reverse implication is proved similarly. □

Proposition 2.2. *For $\{T_n\} \subseteq \mathcal{B}(X)$ assume that $\sup_n \|T_n\| < \infty$. Then $J_{\{T_n\}}(x) = L_{\{T_n\}}(x)$ for every $x \in X$.*

Proof. Note that the inclusion $L_{\{T_n\}}(x) \subseteq J_{\{T_n\}}(x)$ holds trivially. Now suppose $y \in J_{\{T_n\}}(x)$. Then there exist a strictly increasing sequence of positive integers $\{k_n\}$ and a sequence $\{x_n\}$ in X such that $x_n \to x$ and $T_{k_n}x_n \to y$. Put $M = \sup_n \|T_n\|$. Thus we have

$$\begin{aligned} \|T_{k_n}x - y\| &\leq \|T_{k_n}x - T_{k_n}x_n\| + \|T_{k_n}x_n - y\| \\ &\leq M\|x_n - x\| + \|T_{k_n}x_n - y\| \to 0 \end{aligned}$$

as $n \to \infty$. That is $y \in L_{\{T_n\}}(x)$. □

Lemma 2.6. *If $J_{\{T_n\}}(x) = X$ for some non-zero vector $x \in X$, then $J_{\{T_n\}}(\lambda x) = X$ for every $\lambda \in \mathbb{C}$.*

Proof. The case $\lambda \neq 0$ is evident, since it is easy to check that $J_{\{T_n\}}(\lambda x) = \lambda J_{\{T_n\}}(x)$. For the rest of the proof let $\{\lambda_n\}$ be a sequence of non-zero complex numbers converging to 0 and take $y \in J_{\{T_n\}}(x)$. Then by the previous equation

$y \in J_{\{T_n\}}(\lambda_n x)$ for every n. Since $\lambda_n \to 0$, an application of Lemma 2.2 yields that $y \in J_{\{T_n\}}(0)$. Therefore $J_{\{T_n\}}(0) = X$. □

Proposition 2.3. *Let $\{T_n\} \subseteq \mathcal{B}(X)$. Then the set $A = \{x \in X : J_{\{T_n\}}(x) = X\}$ is a closed, connected and $T_n A \subseteq A$.*

Proof. By Lemma 2.2 it is easily inferred that A is closed. To show that A is connected set let $x \in A$ be an arbitrary vector. Then Lemma 2.6, for every $\lambda \in \mathbb{C}$ implies that $J_{\{T_n\}}(0) = J_{\{T_n\}}(\lambda x) = \lambda J_{\{T_n\}}(x) = X$. Thus A is connected. □

Theorem 2.1. *Let $\{T_n\} \subseteq \mathcal{B}(X)$. The following are equivalent.*

(i) The sequence $\{T_n\}$ is hypercyclic;

(ii) The equation $J_{\{T_n\}}(x) = X$ holds for every $x \in X$;

(iii) The set $A = \{x \in X : J_{\{T_n\}}(x) = X\}$ is dense in X;

(iv) The set $A = \{x \in X : J_{\{T_n\}}(x) = X\}$ has non-empty interior.

Proof. To prove that (i) implies (ii) let $x, y \in X$. We know that the set of hypercyclic vectors is G_δ and dense in X (see [1]). Hence there exits a sequence of hypercyclic vectors $\{x_n\}$ such that $x_n \to x$. Subsequently one may find a strictly increasing sequence of positive integers $\{k_n\}$ on which $T_{k_n} x_n \to y$ as $n \to \infty$. Namely $y \in J_{\{T_n\}}(x)$.

The implication $(ii) \Rightarrow (iii)$ is trivial. Moreover applying Lemma 2.2 makes sure that (iii) implies (ii).

Now we show that (iv) implies (ii). Fix $x \in A^\circ$ (interior of A) and let $y \in X$ be an arbitrary. Since $y \in X = J_{\{T_n\}}(x)$ there exist a strictly increasing sequence of positive integers $\{k_n\}$ and a sequence $\{x_n\}$ in X such that $x_n \to x$ and $T_{k_n} x_n \to y$. Without loss of generality we may assume that $x_n \in A$ for every $n \in \mathbb{N}$. Indeed this can be done in sake of the our assumption $x \in A^\circ$.

To prove the implication $(ii) \Rightarrow (i)$, let $\{B_j\}$ be a countable open basis for the relative topology of X. By the Baire category theorem and the hypercyclic vectors description ([1], [12]) it is sufficient to show that the set $\bigcup_{n=1}^{\infty} T_n^{-1}(B_j)$

is dense in X for every j. In fact the set of all hypercyclic vectors for $\{T_n\}$ is equal

$$\bigcap_{j=1}^{\infty}\bigcup_{n=1}^{\infty}T_n^{-1}(B_j).$$

At this moment consider the equivalent definition of $J_{\{T_n\}}(x)$ described in Remark 2.1. Let $x \in X$, the neighborhood U of x and B_j be given. Since $J_{\{T_n\}}(x) = X$, there exists $y \in U$ and $n \in \mathbb{N}$ such that $T_n y \in B_j$ or equivalently $y \in T_n^{-1}(B_j)$. This fact completes the proof since U was chosen arbitrarily. □

Proposition 2.4. *Let $\{T_n\} \subseteq \mathcal{B}(X)$. Then the set $A = \bigcup_{n=1}^{\infty}\{x \in X : J_{T_n}(x) = X\}$ is a closed, connected and $\bigcup_{n=1}^{\infty} T_n A \subseteq A$.*

Proof. Let $x \in A$. There exists $n_0 \in \mathbb{N}$ such that $J_{T_{n_0}}(x) = X$. The T_{n_0}-invariance of $J_{T_{n_0}}(x)$ implies that $J_{T_{n_0}}(T_{n_0}x) = X$. Then $T_{n_0}x \in A$. □

Lemma 2.7. *Let $\{T_n\}$ be a mutually commuting subalgebra of $\mathcal{B}(X)$. Then $\bigcup_{n=1}^{\infty} T_i J_{\{T_n\}}(x) \subseteq J_{\{T_n\}}(T_{n_0}x)$ for some $T_{\{n_0\}} \in \mathcal{B}(X)$.*

Proof. Let $y' \in \bigcup_{n=1}^{\infty} T_i J_{\{T_n\}}(x)$. Then $y' = T_{n_0}y$ for some $T_{n_0} \in \mathcal{B}(X)$ and $y \in J_{\{T_n\}}(x)$. So there exist a strictly increasing sequence of positive integers $\{k_n\}$ and a sequence $\{x_n\}$ in X such that $x_n \to x$ and $T_{k_n}x_n \to y$. Hence $T_{n_0}T_{k_n}x_n \to T_{n_0}y$. Indeed $T_{k_n}T_{n_0}x_n \to y'$ i.e., $y' \in J_{\{T_n\}}(T_{n_0}x)$ since $T_{n_0}x_n \to T_{n_0}x$. □

Proposition 2.5. *Let $\{T_n\} \subseteq \mathcal{B}(X)$ be mutually commuting. Suppose there exists a cyclic vector $x \in X$ for $\{T_n\}$ such that $J_{\{T_n\}}(x)$ has nonempty interior. Then for every nonzero polynomial P and $n \in \mathbb{N}$ the operators $P(T_n)$ has dense range. Moreover the point spectrum $\sigma_p(T_n^*)$ of the adjoint operator of T_n is empty.*

Proof. First assume that X is a complex Banach space. For each $n \in \mathbb{N}$ write $P(T_n) = \alpha(T_n - \lambda_1 I)(T_n - \lambda_2 I)...(T_n - \lambda_k I)$ for some $\alpha, \lambda_i \in \mathbb{C}(i = 1, 2, ..., k)$ where I stands for the identity operator. So it suffices to show that $T_n - \lambda I$ has dense range for every $\lambda \in \mathbb{C}$. Suppose on contrary there exists a nonzero

linear functional x^* such that $x^*(T_n - \lambda I)(x) = 0$ for every $x \in X$. Taking $y \in J^{\circ}_{\{T_n\}}(x)$ is caused to find a strictly increasing sequence of positive integers $\{k_n\}$ and a sequence $\{x_n\} \subseteq X$ such that $x_n \to x$ and $T_{k_n} x_n \to y$. Now by passing to the subsequence $\{k_n\}$ and letting $n \to \infty$ we have $x^*(y) = \lambda x^*(x)$ which is contradiction. □

Theorem 2.2. *Let $\{T_n\} \subseteq \mathcal{B}(X)$ be mutually commuting. Then $\{T_n\}$ is hypercyclic if and only if there exists a cyclic vector $x \in X$ for $\{T_n\}$ such that $J_{\{T_n\}}(x) = X$.*

Proof. We just prove the only if part of assertion. Let $x \in X$ be acyclic vector for $\{T_n\}$, $J_{\{T_n\}}(x) = X$ and let P be an arbitrary nonzero polynomial. By Lemma 2.4 we have $P(T_m)J_{\{T_n\}}(x) \subseteq J_{\{T_n\}}(P(T_m)x)$ for every $m \in \mathbb{N}$. By previous proposition every $P(T_m)$ has dense range. By applying this fact and Lemma 2.4 to the above inclusion we obtain

$$X = \overline{P(T_m)X} \subseteq J_{\{T_n\}}(P(T_m)x).$$

Thus $J_{\{T_n\}}(P(T_m)x) = X$. Now using the fact that x is a cyclic vector follows that there exists a dense subset Y of X on which $J_{\{T_n\}}(y) = X$ for every $y \in Y$. Therefore by Theorem 2.1, $\{T_n\}$ is hypercyclic. □

The following corollary can be readily deduced by the same technique used in the proof of Proposition 2.5.

Corollary 2.1. *Let $\{T_n\} \subseteq \mathcal{B}(X)$. Suppose there exists a vector $x \in X$ such that $J_{\{T_n\}}(x)$ has nonempty interior. Then for every sequence of complex numbers $\{\lambda_n\}$ with $\lambda_n \to 0$ the sequence of bounded linear operators $\{T_n - \lambda_n I\}$ has dense range.*

References

[1] M. R. Azimi, subspace- diskcyclic sequences of linear operators, preprint (2014) 18.

[2] M. R. Azimi, V. Muller, A note on J-sets of linear operators, Rev. R. Acad. Cienc. Exactas Fs. Nat. Ser. A Math. RACSAM 105 (2) (2011) 449453.

[3] F. Bayart, E. Matheron, Dynamics of linear operators, vol. 179 of Cambridge Tracts in Mathematics, Cambridge University Press, Cambridge, 2009.

[4] L. Bernal-Gonzalez, K.-G. Grosse-Erdmann, The hypercyclicity criterion for sequences of operators, Studia Math. 157 (1) (2003) 1732.

[5] J. Bonet, F. Martnez-Gimenez, A. Peris, Linear chaos on Frechet spaces, Internat. J. Bifur. Chaos Appl. Sci. Engrg. 13 (7) (2003) 16491655, dynamical systems and functional equations (Murcia, 2000).

[6] P. S. Bourdon, N. S. Feldman, Somewhere dense orbits are everywhere dense, Indiana Univ. Math. J. 52 (3) (2003) 811819.

[7] G. Costakis, A. Manoussos, J-class weighted shifts on the space of bounded sequences of complex numbers, Integral Equations Operator Theory 62 (2) (2008) 149158.

[8] G. Costakis, A. Manoussos, J-class operators and hypercyclicity, J. Operator Theory 67 (1) (2012) 101119.

[9] G. Godefroy, J. H. Shapiro, Operators with dense, invariant, cyclic vector manifolds, J. Funct. Anal. 98 (2) (1991) 229269.

[10] K.-G. Grosse-Erdmann, Universal families and hypercyclicity operators, Bull. Amer. Math. Soc. 36 (1999) 345381.

[11] F. Leon-Saavedra, V. Muller, Hypercyclic sequences of operators, Studia Math. 175 (1) (2006) 118.

[12] B. F. Madore, R. A. Martnez-Avendano, Subspace hypercyclicity, J. Math. Anal. Appl. 373 (2) (2011) 502511.

[13] Q. Menet, Hypercyclic subspaces and weighted shifts, Adv. Math. 255 (2014) 305337.

[14] A. Montes-Rodrguez, H. N. Salas, Supercyclic subspaces, Bull. London Math. Soc. 35 (6) (2003) 721737.

[15] H. Petersson, A hypercyclicity criterion with applications, J. Math. Anal. Appl. 327 (2) (2007) 14311443.

[16] H. N. Salas, Hypercyclic weighted shifts, Trans. Amer. Math. Soc. 347 (3) (1995) 9931004. 11